



Origination 2/28/2025
Last Approved 4/24/2026
Effective 4/24/2026
Last Revised 2/28/2025
Next Review 4/24/2027

Area Clinical
Services -
Medical
Management
Lines Of All Lines of
Business Business

Air Ambulance Services

PURPOSE:

This Coverage Policy addresses licensed air ambulance transportation services that employ specially designed and equipped vehicles to transport ill or injured patients via air (fixed wing/rotary wing).

POLICY:

Coverage for ambulance services varies across plans. Refer to the customer's benefit plan document for coverage details.

Ambulance transport is considered medically necessary when EITHER of the following conditions is met:

• In connection with a medical emergency AND EITHER of the following criteria are met:

- Transport is to the nearest available provider or facility where the required medical care can be provided
- Transport is from a facility that is not equipped or staffed to treat the patient's specific injury or illness to the nearest appropriate facility where the required medical care can be provided

• **The individual requires specialized medical services during transport and close supervision, or is bed-confined (i.e., cannot get up from bed without assistance, unable to ambulate, and unable to sit in a wheelchair/chair) AND BOTH of the following:**

Either of the following circumstances:

- Transfer is made to the nearest appropriate facility where the required medically necessary and covered diagnostic and/or therapeutic services can be provided to treat the patient's specific injury or illness (e.g., acute care or long-term acute care hospital, acute rehabilitation facility, sub-acute facility, skilled nursing facility, outpatient facility, dialysis center)
- Transfer is made from an inpatient facility, outpatient facility or dialysis center to an individual's home upon completion of required medically necessary and covered diagnostic and/or therapeutic services
- Other means of transport are medically contraindicated or not feasible

Air ambulance transport is considered medically necessary when criteria for ambulance transport are met AND ANY of the following:

- The point of pickup is inaccessible by a ground ambulance
- Great distances or other obstacles are involved in getting the individual to the nearest hospital with appropriate facilities if transport by ground ambulance is not feasible
- Ground ambulance transport would impede timely and appropriate medical care

Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing, transportation, and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

PROCEDURE:

Ambulance services typically involve the assessment, care and transport of an ill or injured patient by specially trained personnel. Patient transportation occurs in safe and monitored environment via specially designed and equipped ground vehicles, aircraft (fixed wing or rotary wing) or boats. Ambulance services must have the necessary permits and licenses in compliance with all the local, state and federal laws and regulations

Professional Societies/Organizations The American College of Emergency Physicians and National Association of EMS Physicians (NAEMSP) published guidelines for the utilization of air medical transport including clinical situations for scene triage, air transport (also known as primary air transport), and inter-

facility transfers. This position statement has been endorsed by the Air Medical Physician Association (AMPA) (Thomson, et al., 2003).

Clinical Indications for scene triage to air transport:

Trauma

A. General and mechanism considerations:

- Trauma Score less than 12, (Glasgow Coma Scale, Systolic Blood, Pressure Respiratory)
- Unstable vital signs (e.g., hypotension or tachypnea)
- Significant trauma in patients less than 12 years old, greater than 55 years old, or pregnant patients
- Multisystem injuries (e.g., long-bone fractures in different extremities; injury to more than two body regions)
- Ejection from vehicle
- Pedestrian or cyclist struck by motor vehicle
- Death in same passenger compartment as patient
- Ground provider perception of significant damage to patient's passenger compartment
- Penetrating trauma to the abdomen, pelvis, chest, neck, or head
- Crush injury to the abdomen, chest, or head
- Fall from significant height

B. Neurologic considerations: Glasgow Coma Scale score less than 10 †

- Deteriorating mental status
- Skull fracture
- Neurologic presentation suggestive of spinal cord injury

C. Thoracic considerations:

- Major chest wall injury (e.g., flail chest)
- Pneumothorax/hemothorax
- Suspected cardiac injury

D. Abdominal/pelvic considerations:

- Significant abdominal pain after blunt trauma
- Presence of a "seatbelt" sign or other abdominal wall contusion
- Obvious rib fracture below the nipple line
- Major pelvic fracture (e.g., unstable pelvic ring disruption, open pelvic fracture, or pelvic fracture with hypotension)

E. Orthopedic/extremity considerations:

- Partial or total amputation of a limb (exclusive of digits)
- Finger/thumb amputation when emergent surgical evaluation (i.e., for replantation consideration) is indicated and rapid surface transport is not available
- Fracture or dislocation with vascular compromise
- Extremity ischemia
- Open long-bone fractures
- Two or more long-bone fractures

F. Major burns:

- Greater than 20% body surface area
- Involvement of face, head, hands, feet, or genitalia
- Inhalational injury
- Electrical or chemical burns
- Burns with associated injuries

G. Patients with near drowning injuries

Clinical Indications for Inter facility Transfers

A. Trauma

- Depending on local hospital capabilities, the Indications listed above under “scene” guidelines may be sufficient indication for air transport; or
- After initial evaluation reveals injuries or potential injuries requiring further evaluation and management beyond the capabilities of the referring hospital.

B. Cardiac

- Acute coronary syndromes with time-critical need for urgent interventional therapy unavailable at the referring center (e.g., cardiac catheterization, intra-aortic balloon pump placement, emergent cardiac surgery)
- Cardiogenic shock
- Cardiac tamponade
- Mechanical cardiac disease (e.g., acute cardiac rupture, decompensating valvular heart disease)

C. Critically ill medical or surgical patients

- Pre-transport cardiac/respiratory arrest
- Requirement for continuous intravenous vasoactive medications or mechanical ventricular assist to maintain stable cardiac output
- Risk for airway deterioration (e.g., angioedema, epiglottitis)
- Acute pulmonary failure and/or requirement for sophisticated pulmonary intensive care (e.g., inverse-ratio ventilation) during transport
- Severe poisoning or overdose requiring specialized toxicology services
- Urgent need for hyperbaric oxygen therapy (e.g., vascular gas embolism, necrotizing infectious)

process, carbon monoxide toxicity)

- Requirement for emergent dialysis
- Gastrointestinal hemorrhages with hemodynamic compromise
- Surgical emergencies such as fasciitis, aortic dissection or aneurysm or extremity ischemia
- Pediatric patients for whom referring facilities cannot provide required evaluation and/or therapy

D. Obstetric

- Reasonable expectation that delivery of infant(s) may require obstetric or neonatal care beyond the capabilities of the referring hospital
- Active premature labor when estimated gestational age is <34 weeks or estimated fetal weight <2,000 grams
- Severe pre-eclampsia or eclampsia
- Third-trimester hemorrhage
- Fetal hydrops
- Maternal medical conditions (e.g., heart disease, drug overdose, metabolic disturbances) exist that may cause premature birth
- Severe predicted fetal heart disease
- Acute abdominal emergencies when estimated gestational age is <34 weeks or estimated fetal weight <2,000 grams

E. Neurological

- Central nervous system hemorrhage
- Spinal cord compression by mass lesion
- Evolving ischemic stroke (i.e., potential candidate for lytic therapy)
- Status epilepticus

F. Neonatal

- Gestational age <30 weeks, body weight <2,000 grams or complicated neonatal course (e.g., perinatal cardiac/respiratory arrest, hemo-dynamic instability, sepsis, meningitis, metabolic derangement, temperature instability)
- Requirement for supplemental oxygen exceeding 60%, continuous positive airway pressure (CPAP), or mechanical ventilation
- Extra pulmonary air leak, interstitial emphysema, or pneumothorax
- Medical emergencies such as seizure activity, congestive heart failure, or disseminated intravascular coagulation
- Surgical emergencies such as diaphragmatic hernia, necrotizing enterocolitis, abdominal wall defects, intussusception, suspected volvulus, or congenital heart defects

G. Transplant

- Patient has met criteria for brain death and air transport is necessary for organ salvage
- Organ and/or organ recipient requires air transport to the transplant center in order to maintain viability of time-critical transplant

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD		No Determination found	
LCD	Palmetto GBA	Ambulance Services (L34549)	7/29/2021

Note: Please review the current Medicare Policy for the most up-to-date information.

References

1. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed Jan 30 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/reports/local-coverage-proposed-lcdsalphabetical-report.aspx?proposedStatus=A&sortBy=title>
2. Centers for Medicare and Medicaid Services (CMS). Medicare Benefit Policy Manual, Chapter 10, Ambulance Services. Accessed on Jan 30, 2025. Available at: <https://www.cms.gov/manuals/Downloads/bp102c10.pdf>
3. Centers for Medicare and Medicaid Services (CMS). Medicare Claims Processing Manual Chapter 15, Ambulance Services. Accessed on Jan 30, 2025. Available at: <http://www.cms.gov/manuals/downloads/clm104c15.pdf>
4. Centers for Medicare and Medicaid Services (CMS). National Coverage Determinations (NCDs) alphabetical index. Accessed Jan 30, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/reports/national-coverage-ncdreport.aspx?chapter=all&sortBy=title>
5. Thomson DP, Thomas SH; American College of Emergency Physicians and National Association of EMS Physicians. Guidelines for air medical dispatch. Position Statement 2002. Prehosp Emerg Care. 2003 Apr-Jun; 7(2):265-71.

All Revision Dates

2/28/2025

COPY